

# On Certain Results for a Subclass of Analytic Functions

Sukhwinder Singh Billing

Department of Applied Sciences  
Baba Banda Singh Bahadur Engineering College  
Fatehgarh Sahib-140 407, Punjab, India  
e-mail: ssbilling@gmail.com

## Abstract

*Using the concept of differential subordination, we extend some results of Al-Abbadi and Darus [1] on analytic univalent functions. Mathematica 7.0 is used to plot the extended regions of the complex plane.*

**Keywords:** *Analytic function, Univalent function, Differential subordination.*

**2000 Mathematical Subject Classification:** Primary 30C80, Secondary 30C45.

## 1 Introduction

Let  $\mathcal{A}$  be the class of all functions  $f$  which are analytic in the open unit disk  $\mathbb{E} = \{z : |z| < 1\}$  and normalized by the conditions that  $f(0) = f'(0) - 1 = 0$ . Thus,  $f \in \mathcal{A}$  has the Taylor series expansion

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k.$$

A function  $f$  is said to be univalent in a domain  $\mathbb{D}$  in the extended complex plane if and only if it is analytic in  $\mathbb{D}$  except for at most one simple pole and  $f(z_1) \neq f(z_2)$  for  $z_1 \neq z_2$  ( $z_1, z_2 \in \mathbb{D}$ ). Let  $\mathcal{S}$  denote the class of all analytic univalent functions  $f$  defined in the unit disk  $\mathbb{E}$  which are normalized by the conditions  $f(0) = f'(0) - 1 = 0$ .

For two functions  $f$  and  $g$  analytic in the open unit disk  $\mathbb{E}$ , we say that  $f$  is subordinate to  $g$  in  $\mathbb{E}$  and write as  $f \prec g$  if there exists a Schwarz function  $w$  analytic in  $\mathbb{E}$  with  $w(0) = 0$  and  $|w(z)| < 1$ ,  $z \in \mathbb{E}$  such that  $f(z) = g(w(z))$ ,  $z \in \mathbb{E}$ . In case the function  $g$  is univalent, the above subordination is equivalent to  $f(0) = g(0)$  and  $f(\mathbb{E}) \subset g(\mathbb{E})$ .

Let  $\psi : \mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}$  be an analytic function,  $p$  be an analytic function in  $\mathbb{E}$ , with  $(p(z), zp'(z)) \in \mathbb{C} \times \mathbb{C}$  for all  $z \in \mathbb{E}$  and  $h$  be univalent in  $\mathbb{E}$ , then a function  $p$  is said to satisfy first order differential subordination if

$$\psi(p(z), zp'(z)) \prec h(z), \quad \psi(p(0), 0) = h(0). \quad (1)$$

A univalent function  $q$  is called a dominant of the differential subordination (1) if  $p(0) = q(0)$  and  $p \prec q$  for all  $p$  satisfying (1). A dominant  $\tilde{q}$  that satisfies  $\tilde{q} \prec q$  for all dominants  $q$  of (1), is said to be the best dominant of (1).

A function  $f \in \mathcal{A}$  is said to be a member of the class  $\mu(\alpha)$  if and only if

$$\left| \frac{4z^2 f'(z)}{[f(z) - f(-z)]^2} - 1 \right| < 1 - \alpha, \quad 0 \leq \alpha < 1, \quad z \in \mathbb{E}.$$

The class  $\mu(\alpha)$  was introduced and studied by Al-Abbadi and Darus [1]. Nunokawa [2] proved the following result.

**Theorem 1.1** *Let  $f \in \mathcal{A}$  satisfy the condition*

$$\left| \frac{z^2 f'(z)}{f^2(z)} - 1 \right| < 1, \quad z \in \mathbb{E},$$

*then  $f$  is univalent in  $\mathbb{E}$ .*

Al-Abbadi and Darus [1] proved the following result.

**Theorem 1.2** *If  $f \in \mathcal{A}$  satisfies*

$$\left| \frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)} \right| < \frac{1 - \alpha}{2 - \alpha}, \quad 0 \leq \alpha < 1, \quad z \in \mathbb{E},$$

*then  $f \in \mu(\alpha)$ .*

As a consequence of above two results, Al-Abbadi and Darus [1] also concluded the following result.

**Theorem 1.3** *If  $f \in \mathcal{A}$  is of the form  $f(z) = z + \sum_{k=2}^{\infty} a_{2k-1} z^{2k-1}$  and satisfies*

$$\left| \frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)} \right| < \frac{1}{2}, \quad z \in \mathbb{E},$$

*then  $f$  is univalent in  $\mathbb{E}$ .*

The main objective of this paper is to extend the region of variability of the operator  $\frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)}$  in above stated results of Al-Abbadi and Darus [1] to get the same conclusion. Mathematica 7.0 is used to show the extended regions of the complex plane.

## 2 Preliminaries

To prove our main result, we shall make use of the following lemma of Miller and Mocanu [3].

**Lemma 2.1** ([3], p.132, Theorem 3.4 h) *Let  $q$  be univalent in  $\mathbb{E}$  and let  $\theta$  and  $\phi$  be analytic in a domain  $\mathbb{D}$  containing  $q(\mathbb{E})$ , with  $\phi(w) \neq 0$ , when  $w \in q(\mathbb{E})$ . Set  $Q(z) = zq'(z)\phi[q(z)]$ ,  $h(z) = \theta[q(z)] + Q(z)$  and suppose that either*

(i)  *$h$  is convex, or*

(ii)  *$Q$  is starlike.*

*In addition, assume that*

(iii)  $\Re \frac{zh'(z)}{Q(z)} > 0, z \in \mathbb{E}$ .

*If  $p$  is analytic in  $\mathbb{E}$ , with  $p(0) = q(0), p(\mathbb{E}) \subset \mathbb{D}$  and*

$$\theta[p(z)] + zp'(z)\phi[p(z)] \prec \theta[q(z)] + zq'(z)\phi[q(z)],$$

*then  $p \prec q$  and  $q$  is the best dominant.*

## 3 Main Result

**Theorem 3.1** *Let  $f \in \mathcal{A}$ ,  $\frac{4z^2 f'(z)}{[f(z) - f(-z)]^2} \neq 0, z \in \mathbb{E}$ , satisfy the differential subordination*

$$\frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)} \prec \frac{\alpha z}{1 + \alpha z}, z \in \mathbb{E},$$

*then*

$$\frac{4z^2 f'(z)}{[f(z) - f(-z)]^2} \prec 1 + \alpha z, 0 < \alpha \leq 1.$$

**Proof.** Let us define the functions  $\theta$  and  $\phi$  as follows:

$$\theta(w) = 0 \quad \text{and} \quad \phi(w) = \frac{1}{w}.$$

Obviously, the functions  $\theta$  and  $\phi$  are analytic in domain  $\mathbb{D} = \mathbb{C} \setminus \{0\}$  and  $\phi(w) \neq 0$  in  $\mathbb{D}$ . Also define the functions  $Q$  and  $h$  as under:

$$Q(z) = zq'(z)\phi(q(z)) = \frac{zq'(z)}{q(z)}$$

and

$$h(z) = \theta(q(z)) + Q(z) = \frac{zq'(z)}{q(z)}.$$

Further, select the functions  $p(z) = \frac{4z^2f'(z)}{[f(z) - f(-z)]^2}$ ,  $f \in \mathcal{A}$  and  $q(z) = 1 + \alpha z$ ,  $0 < \alpha \leq 1$ . Therefore,

$$Q(z) = h(z) = \frac{\alpha z}{1 + \alpha z},$$

and

$$\frac{zQ'(z)}{Q(z)} = \frac{zh'(z)}{Q(z)} = \frac{1}{1 + \alpha z}.$$

It can easily be verified that  $\Re \frac{zQ'(z)}{Q(z)} = \Re \frac{zh'(z)}{Q(z)} > 0$  in  $\mathbb{E}$  for  $0 < \alpha \leq 1$ .

Hence, in view of Lemma 2.1, we obtain  $\frac{4z^2f'(z)}{[f(z) - f(-z)]^2} \prec 1 + \alpha z$ ,  $0 < \alpha \leq 1$  in  $\mathbb{E}$ .

**Remark 3.1** *We, here, make a comparison of the above theorem with the result of Al-Abbadi and Darus [1] stated in Theorem 1.2. We notice that our result extends the region of variability of the differential operator  $\frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)}$  to get the required implication. We compare the results by considering the following particular cases. For  $\alpha = 1$  and  $f$  same as in above theorem, we have:*

$$\begin{aligned} \Re \left( \frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)} \right) &< \frac{1}{2} \\ \Rightarrow \left| \frac{4z^2f'(z)}{[f(z) - f(-z)]^2} - 1 \right| &< 1, \quad z \in \mathbb{E}. \end{aligned} \quad (2)$$

For  $\alpha = 0$ , Theorem 1.2 gives:

$$\begin{aligned} \left| \frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)} \right| &< \frac{1}{2}, \quad z \in \mathbb{E} \\ \Rightarrow \left| \frac{4z^2f'(z)}{[f(z) - f(-z)]^2} - 1 \right| &< 1. \end{aligned} \quad (3)$$

Using Mathematica 7.0, we show our claim pictorially in Figure 2.1. According to the result in (3), the operator  $\frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)}$  takes values in the disk of radius  $\frac{1}{2}$  with center at origin shown by dark shaded region in Figure 2.1 whereas in view of the result in (2), the same operator can take values in the total shaded region (light and dark) to conclude the same result. Thus, the light shaded portion is the claimed extension.




Figure 2.1

**Remark 3.2** In view of the result of Nunokawa [2] stated in Theorem 1.1, if  $\alpha = 1$  and  $f \in \mathcal{A}$  is of the form  $f(z) = z + \sum_{k=2}^{\infty} a_{2k-1}z^{2k-1}$ , we obtain:

$$\Re \left( \frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)} \right) < \frac{1}{2}, \quad z \in \mathbb{E},$$

then  $f$  is univalent in  $\mathbb{E}$ .

We notice that the above result extends the region of variability of the differential operator  $\frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)}$  over the result of Al-Abbadi and Darus [1] stated in Theorem 1.3 to get the required conclusion. By Theorem 1.3, the operator  $\frac{(zf(z))''}{f'(z)} - \frac{2z[f(z) - f(-z)]'}{f(z) - f(-z)}$  takes values in the disk of radius  $\frac{1}{2}$  with center at origin shown by dark shaded region in Figure 2.1 to imply the univalence of  $f \in \mathcal{A}$  whereas according to the above result same operator

can take values in the total shaded region (light and dark) to conclude the same result. Therefore, the light shaded portion shows the extension.

## 4 Open Problem

The technique of differential subordination can similarly be applied to extend the regions of variability of differential operators involving with other classes of analytic univalent functions.

## References

- [1] M. H. Al-Abbadi and M. Darus, *Angular Estimates for Certain Analytic Univalent Functions*, Int. J. Open Problems Complex Analysis, **2**(3)(2010), 211-220.
- [2] M. Nunokawa, *On Some Angular Estimates of Analytic Functions*, Math. Japon., **41**(2)(1995), 447-452.
- [3] S. S. Miller and P. T. Mocanu, *Differential Subordinations : Theory and Applications*, Series on monographs and textbooks in pure and applied mathematics, (No. **225**), Marcel Dekker, New York and Basel, 2000.